

ž ž ž ž

## ü TiO<sub>2</sub>/UV

naserise@tums.ac.ir

ly /y : y / / :

(Ag-TiO<sub>2</sub>)

yy yi ymg/L

i pH

/ y / i y# g/L

Ag-TiO<sub>2</sub>

Ag-TiO<sub>2</sub>

y g/L Ag-TiO<sub>2</sub>

pH:

yy mg/L

fn / E

/ g/L

y g/L

n /

n

Ag-TiO<sub>2</sub>

fl E


! ! ! ! !

UV-E  
(ZnO-E TiO<sub>2</sub>-E)  
(E)  
" "  
"fl E  
TiO<sub>2</sub> " - mg/L  
" flÿnm E (E  
TiO<sub>2</sub> " "  
" Ô  
"fDopingE "  
(E  
"fl E  
P25 TiO<sub>2</sub> " "  
Hombikat (E  
"fl E  
" Ô  
" TiO<sub>2</sub>  
(yE "  
"fl E (Photocatalytic DegradationE  
"fl E "fl E "


Ag-TiO<sub>2</sub> UV  
 pH  
 TiO<sub>2</sub> Ag-TiO<sub>2</sub> Degussa ,25  
 TiO<sub>2</sub>- Ag  
 (Ag/Ti)  
 Scanning Electron Microscope-Energy Dispersive)  
 Seron Technology AIS-2100 (X Ray (SEM-EDX  
 X' Pert MPD  
 (Transmission  
 ZEISS-EM10C Electron Microscopy (TEM)  
 KV (Accelerating voltage)  
 Brunauer- Emmett- Ag-TiO<sub>2</sub>  
 Autosorb 1 Quantachrome Teller (BET  
 nm  
 fl

HCOOH  
 Hole Scavenger  
 Ag-TiO<sub>2</sub>  
 ( Photodeposition  
 Hydrothermal Sol-gel  
 Chemical Photoreduction  
 Vapor Deposition  
 Swamiathan  
 Tryba DB53 DR23  
 ec y  
 Shirzad Siboni  
 UV/TiO<sub>2</sub>  
 pH  
 pH= y min  
 mg/L g/L  
 n / n /  
 Ghanbarian y  
 UV TiO<sub>2</sub>  
 n / LAS

NaOH pH  
N HCl N / L L  
mg/L cm fl WL  
cm ycm  
min  
min rpm mg/L  
µ Ag-TiO<sub>2</sub>  
Perkin-Elmer Lambda  
nm 15-UV/Vis Spectrometer pH=





Ag-TiO<sub>2</sub>



SEM-EDX  
 Ti Ag (at%)  
 wt%  
 Ag/TiO<sub>2</sub>  
 Au  
 TEM SEM  
 (nm)  
 EXRD  
 Ag-TiO<sub>2</sub>  
 P25)  
 BET (TiO<sub>2</sub> Degussa  
 Ag-TiO<sub>2</sub>  
 TiO<sub>2</sub>-P25 doped TiO<sub>2</sub>  
 Ag-TiO<sub>2</sub> m<sup>2</sup>/g TiO<sub>2</sub>-P25  
 / m<sup>2</sup>/g  
 pH pH



Perkin-Elmer  
 Lambda 25-UV/Vis Spectrometer Elmer  
 (DR5000) nm  
 Ag-TiO<sub>2</sub> UV  
 UV  
 pH  
 Ag-TiO<sub>2</sub>/UV  
 UV  
 SPSS16

Ag-doped TiO<sub>2</sub> SEM-EDX




SEM

SEM-EDX




TEM


Sample	Concentration (mg/L)	pH	Time (min)	Removal (%)
Ag-TiO <sub>2</sub>	10	7	60	100
Ag-TiO <sub>2</sub> /UV	10	7	60	100
Ag-TiO <sub>2</sub> /UV	10	9	60	100
Ag-doped TiO <sub>2</sub> /UV	10	7	60	100
Ag-doped TiO <sub>2</sub> /UV	10	9	60	100
Ag-TiO <sub>2</sub>	10	7	30	100
Ag-TiO <sub>2</sub> /UV	10	7	30	100
Ag-TiO <sub>2</sub> /UV	10	9	30	100
Ag-doped TiO <sub>2</sub> /UV	10	7	30	100
Ag-doped TiO <sub>2</sub> /UV	10	9	30	100



(2-Theta-Scale)




$C_0 = \dots \text{mg/L}$        $pH = \dots$   
 $t = \dots \text{min}$        $Ag-TiO_2 = \dots \text{g/L}$  : Ag-TiO<sub>2</sub>/UV




$Ag-TiO_2/UV$        $pH = \dots$        $C_0 = \dots \text{mg/L}$

Ag-doped TiO<sub>2</sub>      Ti      Ag      "

"BET" XRD "TEM" "SEM-EDX"  
 Ag-doped TiO<sub>2</sub>      TEM  
 / nm      Ag-doped TiO<sub>2</sub>  
 TiO<sub>2</sub>  
 SEM-EDX      / nm      nm  
 Ti      Ag (at%)      flwt%



$Ag-TiO_2 = \dots \text{g/L}$        $pH = \dots$  : Ag-TiO<sub>2</sub>/UV



$Ag-TiO_2 = \dots \text{g/L}$  ,  $pH = \dots$        $C_0 = \dots \text{mg/L}$  : Ag-TiO<sub>2</sub>/UV





pH . . . . . / g/L . . . . . "fl - E  
. . . . . " . . . . . " . . . . . y/ g/L  
. . . . . " . . . . . " . . . . .  
. . . . . " . . . . . " . . . . .  
. . . . . - E  
. . . . . Ag-TiO<sub>2</sub> . . . . . fl  
. . . . . fl E . . . . . " . . . . .  
. . . . . y  
. . . . . Yang " . . . . .  
. . . . . Paracetamol  
. . . . . TiO<sub>2</sub>  
. . . . . E  
. . . . . "fl  
. . . . . " . . . . .  
. . . . . y  
. . . . . Guo  
. . . . . TiO<sub>2</sub>  
. . . . . nm  
. . . . . TiO<sub>2</sub>  
. . . . . fl E  
. . . . . nm  
. . . . . fl y Ranjit " . . . . .  
. . . . . " . . . . .  
. . . . . y/ g/L  
. . . . . n  
. . . . . " . . . . .  
. . . . . w . . . . . pH . . . . . E  
. . . . . fl E . . . . . " . . . . . fl  
. . . . . nm  
. . . . . E . . . . . y . . . . . Damodar .( y)  
. . . . . fl

i ymin  
. . . . . " . . . . .  
. . . . . ymin . . . . . fl E  
. . . . . " . . . . .  
. . . . . " . . . . .  
. . . . . ( E . . . . . " . . . . .



- Javadi AH. Photocatalytic degradation of phenol in Aqueous Solutions by Fe(III)-doped TiO<sub>2</sub>/UV Process. *Journal of Health & Environment*. 2011;3(4):369-380 (in Persian).
9. Sa J, Aguera CA, Gross S, Anderson JA. Photocatalytic nitrate reduction over metal modified TiO<sub>2</sub>. *Journal of Applied Catalysis B: Environmental*. 2009;85(3-4):192-200.
10. Van Grieken R, Marugan J, Sordo C, Martinez P, Pablos C. Photocatalytic inactivation of bacteria in water using suspended and immobilized silver-TiO<sub>2</sub>. *Journal of Applied Catalysis B: Environmental*. 2009;93(1-2):112-8.
11. Rengaraj S, Li XZ. Enhanced photocatalytic reduction reaction over Bi<sup>3+</sup>-TiO<sub>2</sub> nanoparticles in presence of formic acid as a hole scavenger. *Chemosphere*. 2007;66(5):930-8.
12. Sobana N, Muruganadham M, Swaminathan M. Nano-Ag particles doped TiO<sub>2</sub> for efficient photodegradation of direct azo dyes. *Journal of Molecular Catalysis A: Chemical*. 2006;258 (1-2): 124-32.
13. Tryba B, Piszcz M, Morawski AW. Photocatalytic and self-cleaning properties of Ag-doped TiO<sub>2</sub>. *Journal of Open Materials Science*. 2010;4(2):5-8.
14. Shirzad Siboni M, Samadi MT, Rahmani A, Khataee A, Bordbar M, Samarghandi MR. Photocatalytic removal of hexavalent chromium and divalent nickel from aqueous solution by UV irradiation in the presence of titanium dioxide nanoparticles. *Iranian Journal of Health and Environment*. 2010;3(3):261-70 (in Persian).
15. Ghanbarian M, Nabizadeh R, Mahvi AH, Nasser S, Naddafi K. Photocatalytic degradation of linear alkyl benzene sulfonate from aqueous solution by TiO<sub>2</sub> nanoparticles. *Iranian Journal of Health and Environment*. 2011;8(4):309-16 (in Persian).
16. Behnajadi MA, Modirshahla N, Shokri M, Rad B. Enhancement photocatalytic activity of TiO<sub>2</sub> nanoparticles by silver doping: Photodeposition versus liquid impregnation methods. *Global Nest Journal*. 2008;10(1):1-7.
17. APHA. AWWA. WEF. APHA. Standard Methods for the Examination of Waters and Wastewaters. 21st ed. Washington, DC: American Public Health Association (APHA); 2005.
18. Shaygani Madad M, Jaleh B, Ashrafe Gh. The effect of thermal operation and particle size on TiO<sub>2</sub> nanoparticle phase changes. *Majlesi Journal of Materials Engineering*. 2011;4(4):51-7.
19. Li Y, White TJ, Lim SH. Low temperature synthesis and microstructural control of titania nano-particles. *Journal of Solid State Chemistry*. 2004;177(4-5):1372-81.
20. Ranjit KT, Viswanathan B. Photocatalytic reduction of nitrite and nitrate ions to ammonia on M/TiO<sub>2</sub> catalysts. *Journal of Photochemistry and Photobiology A: Chemistry*. 1997;108(1):73-8.
21. Yang L, Yu LE, Ray MB. Degradation of paracetamol in aqueous solutions by TiO<sub>2</sub> photocatalysis. *Water Research*. 2008;42(13):3480-8.
22. Mahvi AH, Ghanbarian M, Nasser S, Khairi A. Mineralization and discoloration of textile wastewater by TiO<sub>2</sub> nanoparticles. *Desalination*. 2009;239(1-3):309-16.
23. Sobczynski A, Duczmal L. Photocatalytic destruction of catechol on illuminated titania. *Journal of Reaction Kinetics and Catalysis Letters*. 2004;82(2):213-8.
24. Kashif N, Ouyang F. Parameters effect on heterogeneous photocatalysed degradation of phenol in aqueous dispersion of TiO<sub>2</sub>. *Journal of Environmental Sciences*. 2009;21(4):527-33.
25. Guo Z, Ma R, Li G. Degradation of phenol by nanomaterial TiO<sub>2</sub> in wastewater. *Chemical Engineering Journal*. 2006;119(1):55-9.
26. Damodar RA, Swaminathan T. Performance evaluation of a continuous flow immobilized rotating tube photocatalytic reactor (irtpr) immobilized with TiO<sub>2</sub> catalyst for azo dye degradation. *Chemical Engineering Journal*. 2008;144:59-66.

## Photocatalytic Reduction of Nitrate in Aqueous Solutions using Ag-doped TiO<sub>2</sub>/UV Process

Saeed Parastar<sup>1</sup>, \*Simin Nasser<sup>1</sup>, Amir Hossein Mahvi<sup>1</sup>, Mitra Gholami<sup>1</sup>, Amir Hossein Javadi<sup>2</sup>, Saeedeh Hemmati<sup>1</sup>

<sup>1</sup>Department of Environmental Health Engineering, Faculty of Health, Tehran University of Medical Sciences, Tehran, Iran

<sup>2</sup> Institute of Engineering of Jihad Keshavarzi, Tehran, Iran

Received; 03 March 2011 Accepted; 28 May 2011

### ABSTRACT

**Background and Objectives:** Pollution of water resources to nitrate is an environmental problem in many parts of the world. This problem possibly causes diseases such as methemoglobinemia, lymphatic system cancer and Leukemia. Hence, nitrate control and removal from water resources is necessary. Considering that application of nanomaterials in treatment of environmental pollutants has become an interesting method, in this research use of Ag-doped TiO<sub>2</sub> nanoparticles synthesized through photodeposition produced under UV irradiation was studied for removal of nitrate from aqueous solutions.

**Materials and Methods:** Three nitrate concentrations of 20, 50, and 100 mg/L were considered. In order to determine the effect of Ag-doped TiO<sub>2</sub> nanoparticles on nitrate removal, dosages of 0.1, 0.4, 0.8 and 1.2 g/L nanoparticles were used; pH range of 5-9 was also considered. The effect of Ag-doped TiO<sub>2</sub> nanoparticles both in darkness and under UV irradiation was studied. Moreover, the presence of chloride and sulfate anions on the system removal efficiency was investigated.

**Results:** The optimum performance of nitrate removal (95.5%) was obtained using nitrate concentration of 100 mg/L, in acidic pH and 0.8 g/L Ag-TiO<sub>2</sub>. Increase of nanoparticle dosage up to 0.8 g/L, increased the removal efficiency, but for 1.2 g/L dosage of nanoparticles, the removal efficiency decreased. Maximum reduction performance without nanoparticles, under UV irradiation and under darkness conditions were 32% and 23.3% , respectively. In addition, we found that presence of sulfate and chloride anions in aqueous solution reduced efficiency of nitrate removal.

**Conclusion:** Results of this study showed that Ag-doped TiO<sub>2</sub> nanoparticles may be efficiently used for nitrate removal from aqueous solutions.

**Keywords:** Photocatalytic reduction, Ag-doped TiO<sub>2</sub>, Nitrate, Aqueous solutions

---

\*Corresponding Author: [naserise@tums.ac.ir](mailto:naserise@tums.ac.ir)

Tel: +98 21 88989133, Fax: +98 21 88989133